

PID Curriculum

Pelvic Inflammatory Disease (PID)



1

PID Curriculum

Learning Objectives

Upon completion of this content, the learner will be able to:

1. Describe the epidemiology of PID in the U.S.
2. Describe the pathogenesis of PID.
3. Discuss the clinical manifestations of PID.
4. Identify the clinical criteria used in the diagnosis of PID.
5. List CDC-recommended treatment regimens for PID.
6. Summarize appropriate prevention counseling messages for a patient with PID.
7. Describe public health measures to prevent PID.

2

PID Curriculum

Lessons

- I. Epidemiology: Disease in the U.S.
- II. Pathogenesis
- III. Clinical manifestations
- IV. PID diagnosis
- V. Patient management
- VI. Prevention

3

PID Curriculum

**Lesson I: Epidemiology:
Disease in the U.S.**

4

PID Curriculum

Epidemiology

Pelvic Inflammatory Disease

- Clinical syndrome associated with ascending spread of microorganisms from the vagina or cervix to the endometrium, fallopian tubes, ovaries, and contiguous structures.
- Comprises a spectrum of inflammatory disorders including any combination of endometritis, salpingitis, tubo-ovarian abscess, and pelvic peritonitis.

5

PID Curriculum

Epidemiology

Incidence and Prevalence

- Occurs in approximately 1 million U.S. women annually.
- Annual cost exceeds \$4.2 billion.
- Surveillance and reporting limited by insensitive and nonspecific diagnosis and underreporting.
- Hospitalizations declined through the 80s and early 90s; have remained constant since 1995.
- Reported number of initial visits to physicians' offices has remained unchanged since 1998.

6

PID Curriculum Epidemiology

Normal Cervix with Ectopy



Source: Seattle STD/HIV Prevention Training Center at the University of Washington/
Claire E. Stevens

10

PID Curriculum

Lesson II: Pathogenesis


11


PID Curriculum Pathogenesis


Microbial Etiology

- Most cases of PID are polymicrobial
- Most common pathogens:
 - *N. gonorrhoeae*: recovered from cervix in 30%-80% of women with PID
 - *C. trachomatis*: recovered from cervix in 20%-40% of women with PID
 - *N. gonorrhoeae* and *C. trachomatis* are present in combination in approximately 25%-75% of patients

12








PID Curriculum

Lesson III: Clinical Manifestations

16



PID Curriculum Clinical Manifestations

Sequelae

- Approximately 25% of women with a single episode of PID will experience sequelae, including ectopic pregnancy, infertility, or chronic pelvic pain
- Tubal infertility occurs in 50% of women after three episodes of PID

18

PID Curriculum

Lesson IV: PID Diagnosis

19

PID Curriculum

Diagnosis

Minimum Criteria in the Diagnosis of PID

- Uterine/adnexal tenderness or
- Cervical motion tenderness

20

PID Curriculum

Diagnosis


Additional Criteria to Increase Specificity of Diagnosis

- Temperature >38.3°C (101°F)
- Abnormal cervical or vaginal mucopurulent discharge
- Presence of WBCs on saline wet prep
- Elevated erythrocyte sedimentation rate (ESR)
- Elevated C-reactive protein (CRP)
- Gonorrhea or chlamydia test positive

21

PID Curriculum Diagnosis

Mucopurulent Cervical Discharge (Positive swab test)



Source: Seattle STD/HIV Prevention Training Center at the University of Washington/
Claire E. Stevens and Ronald E. Roddy 22

PID Curriculum Diagnosis

More Specific Criteria

- Endometrial biopsy
- Transvaginal sonography or MRI
- Laparoscopy

23

PID Curriculum

Lesson V: Patient Management

24

PID Curriculum Management

General PID Considerations

- Regimens must provide coverage of *N. gonorrhoeae*, *C. trachomatis*, anaerobes, Gram-negative bacteria, and streptococci
- Treatment should be instituted as early as possible to prevent long term sequelae

25

PID Curriculum Management

Criteria for Hospitalization

- Inability to exclude surgical emergencies
- Pregnancy
- Non-response to oral therapy
- Inability to tolerate an outpatient oral regimen
- Severe illness, nausea and vomiting, high fever or tubo-ovarian abscess
- HIV infection with low CD4 count

26

PID Curriculum Management

Oral Regimens

CDC-recommended oral regimen A

- Ofloxacin 400 mg orally 2 times a day for 14 days, OR
- Levofloxacin 500 mg orally 2 times a day for 14 days

With or Without

- Metronidazole 500 mg orally 2 times a day for 14 days

CDC-recommended oral regimen B

- Ceftriaxone 250 mg IM in a single dose, OR
- Cefoxitin 2 g IM in a single dose and Probenecid 1 g orally in a single dose, OR
- other parenteral third-generation cephalosporin (e.g., Ceftrizoxime, Cefotaxime)
- PLUS Doxycycline 100 mg orally 2 times a day for 14 days

With or Without

- Metronidazole 500 mg orally 2 times a day for 14 days

27

PID Curriculum Management

Follow-Up

- Patients should demonstrate substantial improvement within 72 hours.
- Patients who do not improve usually require hospitalization, additional diagnostic tests, and surgical intervention.
- Some experts recommend rescreening for *C. trachomatis* and *N. gonorrhoeae* 4-6 weeks after completion of therapy in women with documented infection with these pathogens.

28

PID Curriculum Management

Parenteral Regimens

CDC-recommended parenteral regimen A

- Cefotetan 2 g IV every 12 hours, OR
- Cefoxitin 2 g IV every 6 hours
- PLUS doxycycline 100 mg orally or IV every 12 hours

CDC-recommended parenteral regimen B

- Clindamycin 900 mg IV every 8 hours
- PLUS gentamicin loading dose IV of IM (2 mg/kg), followed by maintenance dose (1.5 mg/kg) every 8 hours. Single daily gentamicin dosing may be used.

29

PID Curriculum Management

Parenteral Regimens (continued)

- Continue either of these regimens for at least 24 hours after substantial clinical improvement, then
- Complete a total of 14 days therapy with
 - Doxycycline (100 mg orally twice a day) with regimen A or with
 - Doxycycline or Clindamycin (450 mg orally four times a day), if using regimen B

30

PID Curriculum Management

Alternative Parenteral Regimen

CDC-recommended alternative parenteral regimen

- Ofloxacin 400 mg IV every 12 hours, or
- Levofloxacin 500 mg IV once daily

With or Without

- Metronidazole 500 mg IV every 8 hours, or
- Ampicillin/Sulbactam 3 g IV every 6 hours
- PLUS Doxycycline 100 mg orally or IV every 12 hours

31

PID Curriculum

Lesson VI: Prevention

32

PID Curriculum Prevention

Screening

Recommendations

- To reduce the incidence of PID, screen and treat for chlamydia.
- Annual chlamydia screening for:
 - Sexually active women 25 and under
 - Sexually active women >25 at high risk
- Screen pregnant women in the 1st trimester

33

PID Curriculum Prevention

Partner Management

- Male sex partners of women with PID should be examined and treated if they had sexual contact with the patient during the 60 days preceding the patient's onset of symptoms.

34

PID Curriculum Prevention

Partner Management (continued)

- Male partners of women who have PID caused by *C. trachomatis* or *N. gonorrhoeae* are often asymptomatic.
- Sex partners should be treated empirically with regimens effective against both *C. trachomatis* and *N. gonorrhoeae*, regardless of the apparent etiology of PID or pathogens isolated from the infected woman.

35

PID Curriculum Prevention

Reporting

- Report cases of PID to the local STD program in states where reporting is mandated.
- Gonorrhea and chlamydia are reportable in all states.

36

PID Curriculum Prevention

Patient Counseling and Education

- Nature of the infection
- Transmission
- Risk reduction
 - Assess patient's behavior-change potential
 - Discuss prevention strategies
 - Develop individualized risk-reduction plans

37

PID Curriculum

Case Study

38

PID Curriculum Case Study



History: Jane Wheels

- 24-year-old female who presents reporting lower abdominal pain, cramping, slight fever, and dysuria for 4 days
- P 1001, LMP 2 weeks ago (regular without dysmenorrhea). Uses oral contraceptives (for 2 years).
- Reports gradual onset of symptoms of lower bilateral abdominal discomfort, dysuria (no gross hematuria), abdominal cramping and a slight low-grade fever in the evenings for 4 days. Discomfort has gradually worsened.
- Denies GI disturbances or constipation. Denies vaginal d/c.
- States that she is happily married in a monogamous relationship. Plans another pregnancy in about 6 months. No condom use.
- No history of STDs. Reports occasional yeast infections.
- Douches regularly after menses and intercourse; last douched this morning.

39

PID Curriculum Case Study

Physical Exam

- Vital signs: blood pressure 104/72, pulse 84, temperature 38°C, weight 132
- Neck, chest, breast, heart, and musculoskeletal exam within normal limits. No flank pain on percussion. No CVA tenderness.
- On abdominal exam the patient reports tenderness in the lower quadrants with light palpation. Several small inguinal nodes palpated bilaterally.
- Normal external genitalia without lesions or discharge.
- Speculum exam reveals minimal vaginal discharge with a small amount of visible cervical mucopus.
- Bimanual exam reveals uterine and adnexal tenderness as well as pain with cervical motion. Uterus anterior, midline, smooth, and not enlarged.

40

PID Curriculum Case Study

Questions

1. What should be included in the differential diagnosis?
2. What laboratory tests should be performed or ordered?

41

PID Curriculum Case Study

Laboratory

Results of office diagnostics:

- Urine pregnancy test: negative
- Urine dip stick for nitrates: negative
- Vaginal saline wet mount: vaginal pH was 4.5. Microscopy showed WBCs >10 per HPF, no clue cells, no trichomonads, and the KOH wet mount was negative for budding yeast and hyphae.

3. What is the presumptive diagnosis?
4. How should this patient be managed?
5. What is an appropriate therapeutic regimen?

42

PID Curriculum Case Study

Partner Management

Sex partner: Joseph (spouse)

- First exposure: 4 years ago
- Last exposure: 1 week ago
- Frequency: 2 times per week (vaginal only)



6. How should Joseph be managed?

43

PID Curriculum Case Study

Follow-Up

- On follow up 3 days later, Jane was improved clinically. The culture for gonorrhea was positive. The nucleic acid probe (DNA-probe) for chlamydia was negative.
- Joseph (Jane's husband) came in with Jane at follow-up. He was asymptomatic but did admit to a "one-night stand" while traveling. He was treated. They were offered HIV testing which they accepted.

7. Who is responsible for reporting this case to the local health department?

8. What are appropriate prevention counseling recommendations for this patient?

44
